

Injury patterns in sports-related knee and shoulder injuries in Wrestlers from Haryana Practicing in Government Academies

Dr. Anjli

Assistant Professor, Department of Physiotherapy, Jagannath University, Delhi NCR, Haryana.

Abstract- Wrestling is an Olympic sport for both men and women. It is so ancient since 708 B. C. It is a combat sports that results in various types of injuries that occur in sports. Wrestling and football, which involve frequent player-to-player contact, have a high risk of shoulder injuries. Furthermore, wrestling involves constant contact with the playing surface, exacerbating the condition. According to research from the Center for damage and Policy, football and wrestling are the two sports with the highest risk of major damage to athletes. Aim of this study is to know about injury patterns in sports-related knee and shoulder injuries in wrestlers from Haryana practicing in Government Academies. Data has been collected through off line structured questionnaire of wrestlers and coaches of Government Academies of Haryana State. Sample size for the collection of the data has been collected from the total 551 respondents (wrestlers) on the basis of exclusion and inclusion criteria. In our study population, 40 out of 551 players (7.25%) who experienced shoulder injuries and 77 out of 551 (13.97%) who experienced Knee injuries. In our study, 7.25% of participants reported shoulder injuries.

Keywords: Wrestling, injury patterns, Knee injuries, Shoulder injuries, Government Academies, Haryana.

I. INTRODUCTION

Wrestling is an Olympic sport for both men and women. In this sport, wrestlers can use their legs for offensive and defensive maneuvers. Catch-as-catch-can wrestling, a style where pinning and throwing an opponent to the mat is the primary objective, is where the sport initially gained prominence. The significance of sports in fostering development is increasingly recognized today. According to Singh (2018), "Every human being has a fundamental right of access to physical education and sport, which are essential for the full development of his personality." The "UNESCO 1978 International Charter of Physical Education and Sport" offers this claim.

Records of ancient Olympic wrestling date back to the Greeks in 708 B.C., and wrestling's beginnings can be found in the Sumerians as early as 5000 B.C. (Comstock RD, Knox C, et al. 2006) Since then, wrestling has developed into a variety of forms that are practiced all over the world. The main styles are freestyle, Greco-Roman, and folkstyle. The majority

of athletes in the United States (U.S.) compete in folkstyle, which includes younger athletes in youth programs as well as middle school and high school interscholastic wrestling teams. (Boden B P, Lin W, et al 2002)

Young athletes are more vulnerable to major injuries, such as brain damage, lasting debilitation from fractures, and infrequently, death (Boden B P, Lin W, et al 2002, Boden BP, Jarvis et al 2008, Halloran L 2008, Hewett T E, Pasque C et al 2005, Luckstead E F, Patel D R 2002). Many of these injuries can be classified as significant or catastrophic, meaning that wrestlers will often need emergency care. Depending on the diagnosis, some injuries might require surgery or inpatient care (Boden B P, Lin W, et al 2002, Boden B P, Jarvis et al 2008, Halloran L 2008, Yang J, Peek-Asa C, et al 2007. Additionally, wrestling's particular physical demands and limb postures can result in uncommon but significant injuries that have been documented in the medical literature. (Devgan L L, Gill H S, et al 2006, Giombini A, Di Cesare A et al 2003, Johnson R. 2006, .Ozcakar L, Erol O, et al 2003)

In ancient India, the king ensured that wrestlers had a balanced diet of milk, beats, sweets, sugar, and exquisite desserts despite the ruling elite's frequent mockery of them. Wrestlers used to work out constantly to maintain their health and quality of life. Wrestling was popularized and sanctioned by British authorities in India, allowing its practitioners to join the military. Both Indian warriors and British military personnel create readiness plans. As a result, wrestling became a popular sport in India, and the country is now ranked in the top ten in the world. Not until the 1960s did things start to change. In 1967, India hosted the World Wrestling Championships in New Delhi.

According to research from the Center for damage and Policy, football and wrestling are the two sports with the highest risk of major damage to athletes. (Akbarnejad A, Sayyah M 2012). According to a research by Powell et al., the largest proportion of injuries for which a player missed more than 7 days was in baseball (31.0%) and wrestling (32.6%), with the lowest numbers in field hockey (20.4%) and softball (22.9%) (V Boden BP, Lin W et al 2002) Wrestling injuries to the knee and shoulder joints are the most common. (Boden BP, Jarvis CG et al 2008, Halloran L 2008). So this study is very important to be carried out

II. AIM OF THE STUDY

Aim of this study is to know about injury patterns in sports-related knee and shoulder injuries in wrestlers from Haryana practicing in Government Academies

III. MATERIAL AND METHODS

Data has been collected through off line structured questionnaire of wrestlers and coaches of Government Academies of Haryana State. Sample size for the collection of the data has been collected from the total 551 respondents (wrestlers) on the basis of exclusion and inclusion criteria.

Sample:

Number of Subjects- 551 wrestlers were chosen from all Government Academies of Haryana that were inpatients at their concerned academies were chosen

that were recruited in the Government Academies of Haryana.

- Sample Population- wrestlers
- Sample Design- Survey
- Source of Collection- Government Academies of Haryana State
- Method of Data Collection- The study subjects were wrestlers of Haryana State who were inpatient in government academies. The questionnaires were prepared and filled by both the wrestlers as well as coaches of Haryana for the study according to the objectives as per ethical principles of the research.

The questionnaire was filled with the 30 Government Wrestling Academies with Wrestlers practicing in concerned academies.

Inclusion Criteria

- All subjects should be admitted in Government academies of Haryana for more than one year.
- All subjects should be wrestlers.

Exclusion Criteria

- Subjects should be admitted in Government academies of Haryana for less than one year.
- Subjects that were practicing in private academies.
- Subjects that were practicing in another state rather than Haryana.

Variables of The Study

Variables in research can be categorized into two main types:

Independent variables - Government Wrestling academies of Haryana

Dependent Variables - Injury patterns in sports-related knee and shoulder injuries in Wrestlers

Method of assigning Subjects

Subjects were assigned to survey group randomly.

The study is a Survey design in which self-prepared questionnaire is prepared according

to aims and objectives of the study which was filled by both wrestlers and coaches of Government Academies of Haryana.

Sampling Design

The study is a Survey design in which self-prepared questionnaire is prepared according to aims and objectives of the study which was filled by wrestlers of Government Academies of Haryana.

IV. DATA ANALYSIS

The score was calculated in terms of frequency and % in order to assess the data.

All data was evaluated statistically using the SPSS 10.00 software package.

Five hundred fifty one subjects were used in this study who were wrestlers and coaches between the age group of 13 to 20 years (wrestlers) and coaches from All Government Academies of Haryana State. All subjects were admitted and practicing in Government Academies for one year or more.

Demographic data of the respondents have been collected through a self-structured questionnaire.

Variable s	Opts	Percentag e	Frequenc y
Age	Upto 12 Years	9.25%	51
	13-18 Years	72.59%	400
	19-23 Years	18.14%	100

Variabl es	Opts	Percenta ge	Frequen cy
Gender	Male	64.42%	355
	Female	35.57%	196

Variabl es	Opts	Percenta ge	Frequen cy
Height	Upto 140 cm	0.72%	4
	141-150 cm	9.07%	50
	151-160 cm	19.96%	110
	161-170cm	68.42%	377
	>170 cm	1.81%	10

Variable s	Opts	Percentag e	Frequenc y
Weight	Upto 45 Kg	1.81%	10
	46-60 Kg	36.29%	200
	61-75 Kg	45.37%	250
	76-90 Kg	16.51%	91

Patterns of Knee and Shoulder injuries occurred in Wrestlers in last 5 years

The knee (13.97%) and shoulder (7.25%) were affected during practice and competition which resulted interruption in their practice in their respective academies.

Table 2. Knee Lesions.

Medial Meniscus	5	(5/77) 6.49%
Lateral Meniscus	7	(7/77) 9.09 %
Anterior Cruciate Ligament	8	(8/77) 10.38 %
Posterior Cruciate Ligament	2	(2/77) 2.59 %
Medial Collateral Ligament	5	(5/77) 6.49%
Lateral Collateral Ligament	3	(3/77) 3.89%
Effusion	9	(9/77) 11.68%
Bone Contusion	9	(9/77) 11.68%
Strains	14	(14/77) 18.18%
Cartilage lesions	2	(2/77) 2.59%
Prepatellar bursitis	13	(13/77) 16.88%

Shoulder Lesions (N=40)

Sprain/Tears (muscular and ligamentous)	15	(15/40) 37.5%
Contusion	4	(4/40) 10%
Bursitis	2	(2/40) 5%

Tendinosis	1	(1/40) 2.5%
Fracture	2	(2/40) 5%
Bony Bankart	3	(3/40) 7.5%
Bankart	5	(5/40)12.5%
Hill Sachs'	1	(1/40) 2.5%
SLAP-1	1	(1/40) 2.5%
ALPSA	1	(1/40) 2.5%
Gird	1	(1/40) 2.5%
Perthes	1	(1/40) 2.5%

V. RESULT AND DISCUSSION

Sports injuries occur owing to several risk factors interacting at the same time (Requa RK (1991)). Sports injuries occur when different risk variables interact at the same time (Powell JW (1991)). Injuries vary greatly amongst sports, and particular injury patterns have been identified in different sports. This injury pattern is connected to the varying degrees of risk factors associated with various sports (Taimela S, Kujala UM, et al 1990). Wrestling and football, which involve frequent player-to-player contact, have a high risk of shoulder injuries. Furthermore, wrestling involves constant contact with the playing surface, exacerbating the condition. Even though there is no player-to-player contact in baseball, softball, or volleyball, similar injuries are still common in these sports. Continuous stress on the shoulder as a result of repetitive movements contribute to a high injury rate in volleyball, softball, and baseball (DahmDL and LajamCM (2002)).

Wrestling and football need regular player-to-player contact and so have a high incidence of shoulder injuries. Furthermore, wrestling causes constant contact with the playing surface, exacerbating the condition. Even though there is no player-to-player contact in baseball, softball, or volleyball, similar injuries are still common in these sports. Continuous stress on the shoulder as a result of repetitive movements contribute to a high injury rate in volleyball, softball, and baseball (Hewett TE, Pasque

C et al 2005). In our study population, 40 out of 551 players (7.25%) who experienced shoulder injuries and 77 out of 551 (13.97%) who experienced Knee injuries. In our study, 7.25% of participants reported shoulder injuries. shoulder injuries came in second. Percentage of shoulder injuries has been documented in the range of 3.5-24% of wrestling injuries in the School boys and girls population, and come second only to injuries occurring at the knee (Hewett TE, Pasque C et al 2005, LorishTR, RizzoTD Jr, et al 1992).

The meta-analysis conducted by Hewett et al. found that the exposure-based injury rates for pediatric wrestling injuries ranged from 6.0 to 7.6 injuries per 1,000 athletic exposures. Age, experience, and degree of activity all raised the risk of injury. The head/spine/trunk had the highest frequency of injuries, followed by the upper and lower extremities. Hewett TE, Pasque C et al 2005, Requa RK 1991 studied the most frequently damaged areas were the vertebral column and trunk (34%, 60/176) in four high schools over a two-year period, followed by the lower part of the body (33%, 58/176) and upper extremities (29%, 51/176).

LorishTR, RizzoTD Jr, et al, 1992 documented The sorts of injuries in teenage and preadolescent boys in two significant wrestling competitions and discovered that the primary locations affected were the upper extremities (33%, 73/221 injuries), neck, and back (24%, 53/221). Furthermore, it was noted that older wrestlers and possibly larger wrestlers were more likely to sustain injuries. Wrestling stances frequently expose the head and face to harm. In extremities, the knee, shoulder, and ankle joints are said to be the most frequently involved (Jarret GJ, Orwin JF 1998). The use of legs in FS and hands and arms in GR as per rules may make these wrestlers more susceptible to lower and upper extremity injuries, respectively (Bruce DA, Schut L, et al 1984).

In our study population, higher injuries occurred among wrestlers who practiced freestyle wrestling. Another study by Ngom et al. on children in Dakar discovered that the limbs were more damaged, particularly the upper limbs. The elbow joint sustained the majority of injuries, followed by the

forearm and shoulder. Fractures were most frequently discovered, followed by contusions and dislocations (Ngom G, Mohamed AS, et al, 2017). In our investigation, this was not the case. In general, knee injuries were the most frequent, followed by shoulder injuries. Sprains and strains were the most prevalent injury, followed by contusions. Otero et al. also discovered that injuries to the shoulder were among the top four locations affected, resulting in missed time or surgery (Otero JE, Graves CM, et al 2017).

Every sport has a unique injury profile and level of risk, and injuries differ greatly between sports (Taimela S, Kujala UM, et al 1990). The lower extremities is traditionally the most usually injured part in wrestling, with the knee leading the way, followed by the ankle. These injuries frequently necessitate surgery and are typically season-ending. Prepatellar bursitis is a frequent knee injury that is mostly associated with wrestling. (Strauss RH, Lanese RR. 1982, Mysnyk MC, Wroble RR, et al 1986).

In prospective studies, knee injuries have ranged from 7.6 to 44% of all wrestling injuries (Strauss RH, Lanese RR 1982, Hewett TE, Pasque C, et al. 2005). In the only study where the percentage of knee injuries was less than 10%, Lorish et al. reported injuries sustained by wrestlers between the ages of 6 and 16 during competitions (Lorish TR, Rizzo TD, et al 1992). In our analysis, knee injuries accounted for 13.97% of all reported injuries. These wounds are often rather serious. Over an 11-year span in NCAA wrestling, 65% of injuries requiring surgery were to the knee. In the same study, 21% of injuries resulting in more than a week's absence from competition concerned the knee (Jarret GJ, Orwin JF et al 1998 Barroso BG, Silva JMA, et al 2011).

Barroso et al. discovered that the knee (25.5%) had the most lesions, followed by the shoulder (20%) (Barroso BG, Silva JMA, et al. 2011). In a research by Pasque et al. (Pasque CB, Hewett TE. 2000), the shoulder was the most common site of injury (24%), followed by the knee (17%). Wroble et al. (Wroble RR, Mysnyk MC, et al 1986) found that wrestlers who had previously sustained knee injuries were more likely to sustain another injury.

In their study, Barroso et al. discovered that 77% of all injuries were acute (new injuries), 10% were recurrent, 2% were unresolved injuries from the previous year, and 1% were caused by a recent worsening of an unresolved injury. In our analysis, 28.16% of knee injuries were recurring, compared to 71.83% of new injuries.

Sprains account for between thirty and sixty-five percent of all knee injuries. Meniscal injuries are also common, with a reasonably high proportion of lateral to medial tears (Hewett TE, Pasque C, et al 2005). Wroble et al. found that the most common knee injuries were prepatellar bursitis, lateral and medial collateral ligament sprains, and meniscal tears. The highest rates of sprains (23.37%) and strains (18.18%) of muscles and ligaments were also seen in our study.

Sprains account for between thirty and sixty-five percent of all knee injuries. Meniscal injuries are also common, with a reasonably high proportion of lateral to medial tears (Hewett TE, Pasque C, et al 2005). Wroble et al. found that the most common knee injuries were prepatellar bursitis, lateral and medial collateral ligament sprains, and meniscal tears. The highest rates of sprains (23.37%) and strains (18.18%) of muscles and ligaments were also seen in our study.

There are no widely recognized standards for determining how serious a sports injury is (Jarret GJ, Orwin JF, 1998). The requirement for surgical therapy was employed as a severity criterion by Barroso et al. (Barroso BG, Silva JMA, et al 2011). In their study on wrestling players, 13 (9%) lesions were operated on, the majority of which were in the knee. (Agel J, Ransone J, 2003) and Wroble et al. (Wroble RR, Mysnyk MC, et al 1983) found that the majority of surgically treated lesions in wrestling athletes occurred in the knee. In our study, ten wrestlers required operational intervention, nine of which were for knee injuries.

According to Wroble et al., there were 11.5 knee injuries per 100 wrestlers per year that required a week or more of time off (Wroble RR, Mysnyk MC, et al 1983). According to Pasque et al. (Pasque CB,

Hewett TE. 2000), the average amount of time lost due to an injury was five days, with a range of one to thirty-nine days. In our analysis, there were 18.11 knee injuries per 100 wrestlers annually that resulted in a week or more of missed time in competitions as well as in practice sessions also.

VI. LIMITATIONS/WEAKNESS OF OUR STUDY

This study is limited to Haryana only. Hence, we kept the study very simple. In spite of that we observed the following during the course of this study:

- 1) Ignorance of study population and trainers who practice in Private academies persist in treating injuries according to traditional methods.
- 2) Ignorance about importance of reporting injuries/avoidance to continue practicing.
- 3) Lack of proper follow up to continue playing in competitions.

VII. CONCLUSION

The best way to minimize injuries to the athletes is by developing a well-designed and operational injury prevention program by consistently and professionally evaluating the injury patterns and ensure availability of physiotherapists during competitions and practice so that proper treatment could be given to athletes which can boost their play life and performance in their athletic career. Even though not all injuries can be avoided, the best way to minimize them is by well defined injury prevention strategies. The aim of our study was to understand the variables leading to injuries in the Indian wrestlers.

REFERENCES

1. Agel J, Ransone J, Dick R, Oppliger R, Marshall SW. Descriptive epidemiology of collegiate men's wrestling injuries: National Collegiate Athletic Association Injury Surveillance System, 1988-1989 through 2003-2004. *J Athl Train.* 2007;42(2):303-10.
2. Baker BE, Peckham AC, Puppato F, Sanborn JC. Review of meniscal injury and associated sports. *Am J Sports Med.* 1985;13(1):1-4. doi: 10.1177/036354658501300101.
3. Barroso BG, Silva JMA, Garcia AC, Ramos NCO, Martinelli MO, Resende VR, et al. Musculoskeletal injuries in wrestling athletes. *Acta Ortop Bras.* 2011;19(2):98-101.
4. Barroso BG, Silva JMA, Garcia AC, Ramos NCO, Martinelli MO, Resende VR, et al. Musculoskeletal injuries in wrestling athletes. *Acta Ortop Bras.* 2011;19(2):98-101.
5. Boden BP, Jarvis CG. Spinal injuries in sports. *Neurol Clin.* 2008;26:63-78. viii. doi: 10.1016/j.ncl.2007.12.005.
6. Boden BP, Lin W, Young M, et al. Catastrophic injuries in wrestlers. *Am J Sports Med.* 2002;30:791-5. doi: 10.1177/03635465020300060601.
7. Bruce DA, Schut L, Sutton LN (1984) Brain and cervical spine injuries occurring during organized sports activities in children and adolescents. *Prim Care* 11: 175-194.
8. Comstock RD, Knox C, Yard E, et al. Sports-related injuries among high school athletes--United States, 2005-06 school year. *MMWR Morb Mortal Wkly Rep.* 2006 Sep 29;55:1037-40.
9. Dahm DL and Lajam CM (2002) Shoulder instability in the female athlete. *Op Tech Sports Med* 10: 5-9.
10. Devgan LL, Gill HS, Faustin C, et al. Posterior dislocation in a voluntary subluxator: a case report. *Med Sci Sports Exerc.* 2006;38:613-617. doi: 10.1249/01.mss.0000210210.40694.df.
11. Giombini A, Di Cesare A, Sardella F, et al. Myositis ossificans as a complication of a muscle tendon junction strain of long head of biceps. A case report. *J Sports Med Phys Fitness.* 2003;43:75-7.
12. Halloran L. Wrestling injuries. *Orthop Nurs.* 2008;27:189-92. doi: 10.1097/01.NOR.0000320548.20611.16.
13. Hewett TE, Pasque C, Heyl R, et al. Wrestling injuries. *Med Sport Sci.* 2005;48:152-78. doi: 10.1159/000084288.
14. Hewett TE, Pasque C, Heyl R, Wroble R (2005) Wrestling Injuries. *Med Sport Sci* 48: 152-178. Requa R and Garrick JG (1981) Injuries in Interscholastic Wrestling. *Phys Sports med* 9: 44-51.

15. Hewett TE, Pasque C, Heyl R, Wroble R. Wrestling injuries. *Med Sport Sci.* 2005;48:152–78. doi: 10.1159/000084288.
16. Jarret GJ, Orwin JF, Dick RW (1998) Injuries in Collegiate wrestling *AmJ Sports Med* 26: 674-680.
17. Jarret GJ, Orwin JF, Dick RW. Injuries in collegiate wrestling. *Am J Sports Med.* 1998;26:674–80. doi: 10.1177/03635465980260051301.
18. Johnson R. Abdominal wall injuries: rectus abdominis strains, oblique strains, rectus sheath hematoma. *Curr Sports Med Rep.* 2006;5:99–103. doi: 10.1007/s11932-006-0038-8.
19. kbarnejad A, Sayyah M. Frequency of sports trauma in elite national level greco-roman wrestling competitions. *Arch Trauma Res.* 2012;1(2):51–3. doi: 10.5812/at.6866. [DOI] [PMC free article] [PubMed] [Google Scholar]
20. Kordi R, Ziaee V, Rostami M, Wallace WA. Sports injuries and health problems among wrestlers in Tehran. *J Pak Med Assoc.* 2012;62(3):204–8. [PubMed] [Google Scholar]
21. Lorish TR, Rizzo TD, Ilstrup DM, Scott SG. Injuries in adolescent and preadolescent boys at two large wrestling tournaments. *Am J Sports Med.* 1992;20(2):199–202. doi: 10.1177/036354659202000218. [DOI] [PubMed] [Google Scholar]
22. LorishTR, RizzoTD Jr, Ilstrup DM, Scott SG (1992) Injuries in adolescent and preadolescent boys at two large wrestling tournaments. *AmJ Sports Med* 20: 199-202
23. Luckstead EF, Patel DR. Catastrophic pediatric sports injuries. *Pediatr Clin North Am.* 2002;49:581–91. doi: 10.1016/s0031-3955(02)00006-8. [DOI] [PubMed] [Google Scholar]
24. Mysnyk MC, Wroble RR, Foster DT, Albright JP. Prepatellar bursitis in wrestlers. *Am J Sports Med.* 1986;14(1):46–54. doi: 10.1177/036354658601400109. [DOI] [PubMed] [Google Scholar]
25. Ngom G, Mohamed AS, Mbaye PA, Fall M, Ndour O, et al. (2017) Wrestling game injuries among children in Dakar: a report on 172 cases. *Pan African Medical Journal* 26: 150.
26. Oppliger RA, Landry GL, Foster SW, et al. Wisconsin minimum weight program reduces weight-cutting practices of high school wrestlers. *Clin J Sport Med.* 1998;8:26–31. doi: 10.1097/00042752-199801000-00007. [DOI] [PubMed] [Google Scholar]
27. Otero JE, Graves CM, Bollier MJ (2017) Injuries in collegiate wrestlers at an Elite Division I NCAA wrestling Program: An epidemiological study. *Iowa Orthop J* 37: 65-70.
28. Ozcakar L, Erol O, Kara M, et al. Accessory nerve injury during amateur wrestling: silent but not overlooked. *Br J Sports Med.* 2003;37:372. doi: 10.1136/bjism.37.4.372. [DOI] [PMC free article] [PubMed] [Google Scholar]
29. Pasque CB, Hewett TE. A prospective study of high school wrestling injuries. *Am J Sports Med.* 2000;28(4):509–15. doi: 10.1177/03635465000280041101. [DOI] [PubMed] [Google Scholar]
30. Pasque CB, Hewett TE. A prospective study of high school wrestling injuries. *Am J Sports Med.* 2000;28(4):509–15. doi: 10.1177/03635465000280041101. [DOI] [PubMed] [Google Scholar]
31. Pearsall AWt, Russell GV., Jr Ipsilateral clavicle fracture, sternoclavicular joint subluxation, and long thoracic nerve injury: an unusual constellation of injuries sustained during wrestling. *Am J Sports Med.* 2000;28:904–8. doi: 10.1177/03635465000280062301. [DOI] [PubMed] [Google Scholar]
32. Plaga BR, Looby P, Feldhaus SJ, et al. Axillary artery injury secondary to inferior shoulder dislocation. *J Emerg Med.* doi: 10.1016/j.jemermed.2008.01.018. [DOI] [PubMed] [Google Scholar]
33. Powell JW (1991) Epidemiologic research for injury prevention programs in sports. In: Mueller FO, Ryan AJ, Prevention of Athletic Injuries: The Role of the Sports Medicine Team. Philadelphia, PA: FA Davis: 1991: 11-25.
34. Powell JW, Barber-Foss KD. Injury patterns in selected high school sports: A review of the 1995–1997 seasons. *J Athl Train.* 1999;34:277–84. [PMC free article] [PubMed] [Google Scholar]
35. Powell JW, Barber-Foss KD. Injury patterns in selected high school sports: a review of the 1995-1997 seasons. *J Athl Train.* 1999;34(3):277–84. [PMC free article] [PubMed] [Google Scholar]

36. Powell JW, Barber-Foss KD. Traumatic brain injury in high school athletes. *JAMA*. 1999;282(10):958–63. doi: 10.1001/jama.282.10.958. [DOI] [PubMed] [Google Scholar]
37. Requa RK (1991) The scope of the problem: the impact of sport-related injuries. In: Proceedings of Sports Injuries in Youth Surveillance Strategies, Bethesda, MD, 7-8 April 1991 Bethesda, MD. National Institutes of Health: 1991.
38. Roots and History of Olympic Wrestling. FILA Wrestling- Federation Internationale des Lutttes Associees; 2009.
39. Sikka RS, Neault M, Guanche CA. An avulsion of the subscapularis in a skeletally immature patient. *Am J Sports Med*. 2004;32:246–9. doi: 10.1177/0363546503258906. [DOI] [PubMed] [Google Scholar]
40. Sports Business Research Network. High School Participation: Boys Athletics; Number of Part./Schools with Prog., By Sport. 2008. Market Research Results. [Google Scholar]
41. Sports Business Research Network. Wrestling Market Research Results: Participation. 2008.
42. Strauss RH, Lanese RR. Injuries among wrestlers in school and college tournaments. *JAMA*. 1982;248(16):2016–9. [PubMed] [Google Scholar]
43. Taimela S, Kujala UM, Ostemmn K (1990) Intrinsic risk factors and athletic injuries. *Sports Med* 9: 205-215. DahmDL and LajamCM (2002) Shoulder inst
44. Taimela S, Kujala UM, Osterman K. Intrinsic risk factors and athletic injuries. *Sports Med*. 1990;9(4):205–15. doi: 10.2165/00007256-199009040-00002. [DOI] [PubMed] [Google Scholar]
45. Wroble RR, Mysnyk MC, Foster DT, Albright JP. Patterns of knee injuries in wrestling: a six year study. *Am J Sports Med*. 1986;14(1):55–66. doi: 10.1177/036354658601400110. [DOI] [PubMed] [Google Scholar]
46. Yang J, Peek-Asa C, Allareddy V, et al. Patient and hospital characteristics associated with length of stay and hospital charges for pediatric sports-related injury hospitalizations in the United States, 2000–2003. *Pediatrics*. 2007;119:e813–20. doi: 10.1542/peds.2006-2140. [DOI] [PubMed] [Google Scholar]